arXiv:1404.6281v2 [cs.IT] 18 May 2014

EXPLICIT FACTORIZATION OF $x^n - 1 \in \mathbb{F}_q[x]$

F. E. BROCHERO MARTÍNEZ, C. R. GIRALDO VERGARA,
AND L. BATISTA DE OLIVEIRA

Abstract. Let $\mathbb{F}_q$ be a finite field and n a positive integer. In this article, we prove that, under some conditions on q and n, the polynomial $x^n - 1 \in \mathbb{F}_q[x]$ can be split into irreducible binomials $x^t - a$ and an explicit factorization into irreducible factors is given.

Finally, weakening one of our hypothesis, we also obtain factors of the form $x^{2t} - ax^t + b$ and explicit splitting of $x^n - 1$ into irreducible factors.

1. Introduction

The factorization of a polynomial over a finite field has theoretical and practical important consequences in a wide variety of technological situations, including efficient and secure communications, error-correcting codes, deterministic simulations of random processes and digital tracking systems (see for instance [4]); in particular, each irreducible factor of $x^n - 1$ in $\mathbb{F}_q[x]$ determines a cyclic code of length n over $\mathbb{F}_q$ (see [6]). In fact, each cyclic code of length n can be represented as an ideal of the ring $\mathcal{R}_n = \mathbb{F}_q[x]/\langle x^n - 1\rangle$, and each ideal of $\mathcal{R}_n$ is generated by a unique factor of $x^n - 1$.

It is well known that $x^n - 1 = \prod_{d|n} \Phi_d(x)$, where $\Phi_d(x)$ denotes the d-th cyclotomic polynomial (see [5] Theorem 2.45). It follows that the factorization of $x^n - 1$ depends strongly on the factorization of the cyclotomic polynomial. In general, a "generic efficient algorithm" to split $\Phi_d(x)$ in $\mathbb{F}_q[x]$ for arbitrary d and q is an open problem and just some particular cases are known. The complete factorization of $\Phi_{2^k}(x) = x^{2^k} + 1$ over $\mathbb{F}_q[x]$ with $q \equiv 1 \pmod 4$ is a classical result, and in the case where $q \equiv 3 \pmod 4$ the factorization was obtained way by Meyn in [8] within an elementary; in [7] Fitzgerald and Yucas studied how to find the explicit factorization of $\Phi_{2^k r}(x)$, where r is an odd prime and $q \equiv \pm 1 \pmod r$, in particular, they obtained the explicit factorization in the case where $r = 3$. In [9], Wang and Wang give a complete explicit factorization of $\Phi_{2^n \cdot 5}(x)$. Finally, in [3] Chen, Li and Tuerhong found the explicit factorization of $x^{2^m p^n} - 1$ over $\mathbb{F}_q$, where p is an odd prime with $q \equiv 1 \pmod p$.

In this paper, we characterize every irreducible factor of $x^n - 1$ over $\mathbb{F}_q$ assuming that every prime divisor of n is a divisor of $q-1$, and we give explicit expressions for the factorization of $x^n - 1$, so generalizing the results listed above. Observe that, in this case, all factors are binomials or trinomials, thus every irreducible factor is a sparse polynomial (polynomial with few nonzero term) and this type of polynomials

Date: May 20, 2014.
2010 *Mathematics Subject Classification.* 12E05(primary) and 94B05(secondary).
Key words and phrases. Irreducible factor, Cyclotomic Polynomial, Cyclic Codes.

has important applications, for instance, for an efficient hardware implementation of feedback shift registers (see [1]).

2. Preliminaries

Throughout this paper, $\mathbb{F}_q$ will denote a finite field of order q, where q is a power of a prime, θ is a generator of the cyclic group $\mathbb{F}_q^*$; for each $a \in \mathbb{F}_q^*$, $\operatorname{ord}_q a$ will denote the minimum positive integer k such that $a^k = 1$. For each prime p and each integer m, $\nu_p(m)$ will denote the maximum power of p that divide m and $\operatorname{rad}(m)$ denotes the radical of m, i.e., if $m = p_1^{\alpha_1} p_2^{\alpha_2} \cdots p_l^{\alpha_l}$ is the factorization of m in prime factors, then $\operatorname{rad}(m) = p_1 p_2 \cdots p_l$.

It is easy to see that if n divides $q-1$ then $\mathbb{F}_q^*$ contains a primitive n-th root of unity $\zeta_n = \theta^{(q-1)/n} \in \mathbb{F}_q^*$, and the polynomial $x^n - 1$ splits into linear factor of the form $x - \zeta_n^j$, where $j = 1, 2, \ldots, n$, thus each factor is a binomial. A natural question is when $x^n - 1$ can be factorized in a way that each irreducible factor is a binomial of the form $x^t - a$. This question can be divide in three parts:

- Determine when $x^t - a$ is an irreducible polynomial.
- Determine when $x^t - a$ is a factor of $x^n - 1$.
- Find conditions over n and q such that every irreducible factor is a binomial.

Observe that the first part is a classical remarkable result:

Lemma 2.1 (Theorem 3.75 in [5])**.** *Let $t \geq 2$ be an integer and $a \in \mathbb{F}_q^*$. Then the binomial $x^t - a$ is irreducible in $\mathbb{F}_q[x]$ if and only if the following three conditions are satisfied:*

(1) $\operatorname{rad}(t)$ *divides* $\operatorname{ord}_q a$.
(2) $\gcd(t, \frac{q-1}{\operatorname{ord}_q a}) = 1$.
(3) if $4|t$ *then* $q \equiv 1 \pmod 4$.

Remark 2.2. *If $t = 4s$, $q \equiv 3 \pmod 4$ and a is not a square in $\mathbb{F}_q$, then $x^t - a = (x^{2s} - bx^s + c)(x^{2s} + bx^s + c)$, where b and c are elements of $\mathbb{F}_q$ such that $c^2 = -a$ and $b^2 = 2c$.*

For the second question, we can find elementary conditions over t and a:

Lemma 2.3. *The binomial $x^t - a \in \mathbb{F}_q[x]$ is a factor of $x^n - 1$ if and only if $t|n$ and $\operatorname{ord}_q a | \gcd(q-1, \frac{n}{t})$.*

Proof: Using the division algorithm we know that $n = tm + r$ where $0 \leq r < t$. Thus

$$x^n - 1 = (x^t)^m x^r - 1 \equiv a^m x^r - 1 \mod (x^t - a),$$

so, it follows that $x^t - a$ divides $x^n - 1$ if and only if it also divides $a^m x^r - 1$. Since $\deg(a^m x^r - 1) < \deg(x^t - a)$, this happens if and only if $r = 0$ and $a^m = 1$. Therefore t is a divisor of n and $\operatorname{ord}_q a | m = \frac{n}{t}$. Since $\operatorname{ord}_q a | (q-1)$, we conclude that $\operatorname{ord}_q a | \gcd(q-1, \frac{n}{t})$. □

The third part, that will be solved in the next section, depends on some known elementary properties of the cyclotomic polynomial that we present, without proof, in the following proposition.

Proposition 2.4. *Suppose that m is a positive integer and take p a prime such that $\gcd(pm, q) = 1$. Then in a finite field $\mathbb{F}_q$ the following properties of the cyclotomic polynomial are valid:*

(a) $\Phi_{mp}(x) = \dfrac{\Phi_m(x^p)}{\Phi_m(x)}$ *if* $p \nmid m$;
(b) $\Phi_{mp}(x) = \Phi_m(x^p)$ *if* $p \mid m$;
(c) $\Phi_{2m}(x) = \Phi_m(-x)$ *if* $m \geq 3$ *and* m *is an odd number.*

3. Factors of $x^n - 1$

In this section, we prove, imposing some conditions on n and q, that every irreducible factor of $x^n - 1$ is a binomial, and then, removing one of those conditions, we prove that every irreducible factor is a binomial or a trinomial.

The following proposition show that these condition are indeed necessary.

Proposition 3.1. *If every irreducible factor of* $x^n - 1 \in \mathbb{F}_q[x]$ *is a binomial of the form* $x^t - a$, *then* $\operatorname{rad}(n) | (q-1)$ *and either* $8 \nmid n$ *or* $q \not\equiv 3 \pmod 4$.

Proof: Suppose that p is a prime such that $p|n$. Since $\Phi_p(x)$ divides $x^p - 1$ and $x^p - 1$ divide $x^n - 1$, it follows that every irreducible factor of $\Phi_p(x)$ is a binomial of the form $x^t - a$. Since $x^t - a | (x^p - 1)$, by Lemma 2.3 we have that $t|p$, so $t = 1$ or p. In addition, by Theorem 2.47 (ii) in [5], every irreducible factor of $\Phi_p(x)$ has degree $\operatorname{ord}_p q$ which is a divisor of $\varphi(p) = p - 1$. It follows that $t = \operatorname{ord}_p q$ is a common divisor of p and $p - 1$, therefore $t = 1$. It follows that $\operatorname{ord}_p q = 1$ or equivalently $p|(q-1)$.

On the other hand, if $8|n$ then $x^8 - 1$ divides $x^n - 1$ and if $q \equiv 3 \pmod 4$ then -1 is not a square in $\mathbb{F}_q$. In the case that 2 is a square in $\mathbb{F}_q$, we have that

$$x^8 - 1 = (x-1)(x+1)(x^2+1)(x^2+bx+1)(x^2-bx+1),$$

where $b^2 = 2$. The case that 2 is not a square in $\mathbb{F}_q$, then -2 is a square, therefore

$$x^8 - 1 = (x-1)(x+1)(x^2+1)(x^2+bx-1)(x^2-bx-1),$$

where $b^2 = -2$. So there exist irreducible factors of $x^n - 1$ which are not binomials. □

The following result shows that these conditions are in fact sufficient conditions.

Theorem 3.2. *Let* $\mathbb{F}_q$ *be a finite field and* n *a positive integer such that*

(1) $q \not\equiv 3 \pmod 4$ *or* $8 \nmid n$.
(2) $rad(n)|(q-1)$.

Then every irreducible factor of $x^n - 1$ *in* $\mathbb{F}_q[x]$ *is of the form* $x^t - a$, *where* $t \in \mathbb{N}$ *and* $a \in \mathbb{F}_q$ *satisfy the condition of Lemmas 2.1 and 2.3.*

Proof: We proceed by induction over $\Omega(n)$, the total number of prime power factors of n (i.e., $\Omega(n) = \alpha_1 + \cdots + \alpha_k$ where $n = p_1^{\alpha_1} \cdots p_k^{\alpha_k}$).

Observe that if $n|(q-1)$, then there exists an element $\zeta_n \in \mathbb{F}_q$, a primitive n-th root of unity in $\mathbb{F}_q$, and in this case

$$x^n - 1 = \prod_{j=0}^{n-1} (x - \zeta_n^j),$$

so, the theorem is clearly true. In particular, if $\Omega(n) = 1$ then $n = \operatorname{rad}(n)$ is a prime that divides $q - 1$, therefore the first step of the induction is true.

Suppose now that the theorem is true for all n such that $\operatorname{rad}(n)|(q-1)$ and $\Omega(n) \leq N \in \mathbb{N}$ for some $N \geq 1$.

Let n be an integer such that $rad(n)|(q-1)$ and $\Omega(n) = N+1$. Observe that

$$x^n - 1 = \prod_{d|n} \Phi_d(x) = \Phi_n(x) \cdot \prod_{\substack{d|n \\ d \neq n}} \Phi_d(x),$$

and for all $d|n$, $d \neq n$ we have that $\Phi_d(x)|(x^d - 1)$ Then, by induction hypothesis we have that every irreducible factor of $\Phi_d(x)$ is of the form $x^t - a$, so we only need to verify the condition of Lemma 2.3. In fact, since t divides $\frac{d}{gcd(d,q-1)}$ then it divides $\frac{n}{gcd(n,q-1)}$. In addition, $\operatorname{ord}_q a | \gcd(q-1, \frac{d}{t}) | \gcd(q-1, \frac{n}{t})$.

Therefore, it is enough to analyse the factors of $\Phi_n(x)$. Since the theorem is true when $n|(q-1)$, we can suppose without loss of generality that there exists a prime p divisor of n, such that $\nu_p(n) > \nu_p(q-1) \geq \nu_p(\operatorname{rad}(n)) = 1$. So we can assume that $n = pm$ with $\nu_p(m) \geq \nu_p(q-1) \geq 1$.

At this point, we consider two cases:

(1) $p \neq 2$ or $q \not\equiv 3 \pmod 4$
(2) $p = 2$, $q \equiv 3 \pmod 4$ and $\nu_{p'}(n) \leq \nu_{p'}(q-1)$ for each odd prime factor p' of n .

In the first case, by Proposition 2.4 we have

$$\Phi_n(x) = \Phi_{pm}(x) = \Phi_m(x^p).$$

Since $\Omega(m) < \Omega(n)$ it follows by induction hypothesis that every irreducible factor of $\Phi_m(x)$ is of the form $x^t - a$ and so it satisfies the condition of Lemmas 2.1 and 2.3. In addition, $x^{pt} - a$ is a factor, not necessarily irreducible, of $\Phi_m(x^p)$.

Observe that $\gcd(tp, \frac{q-1}{\operatorname{ord}_q a}) = 1$ or p. When the greatest common divisor is 1, we have that $\nu_p(\operatorname{ord}_q a) = \nu_p(q-1) \geq 1$, then $p | \operatorname{ord}_q a$, and therefore $\operatorname{rad}(pt)$ divides $\operatorname{ord}_q a$, so it follows that the first two conditions of Lemma 2.1 are satisfied. In addition, if $q \equiv 3 \pmod 4$ it follows that $p \neq 2$, and by induction hypothesis we know that $4 \nmid t$, therefore $4 \nmid pt$, consequently the third condition of Lemma 2.1 is also satisfied and then $x^{pt} - a$ is irreducible. The condition of Lemma 2.3 are satisfied, because we have already proved that $x^{pt} - a$ is a factor of $x^n - 1$.

When the greatest common divisor is p, we have that $\nu_p(q-1) > \nu_p(\operatorname{ord}_q a)$ and $p \nmid t$. Then $\operatorname{ord}_q a | \frac{q-1}{p}$, or equivalently, a is a root of the polynomial $P(x) = x^{\frac{q-1}{p}} - 1$. Since the roots of this polynomial are $\{\theta^{jp} | j = 1, 2, \ldots, \frac{q-1}{p}\}$, so there exists $b \in \mathbb{F}_q^*$ such that $a = b^p$, and therefore

$$x^{pt} - a = x^{pt} - b^p = \prod_{j=1}^{p} (x^t - \zeta_p^j b),$$

where $\zeta_p \in \mathbb{F}_q$ is a primitive p-th root of unity. Note that

$$\operatorname{ord}_q(\zeta_p^j b) = \operatorname{lcm}(\operatorname{ord}_p \zeta_p^j, \operatorname{ord}_p b) = \operatorname{lcm}(p, p \cdot \operatorname{ord}_p a) = p \cdot \operatorname{ord}_q a,$$

so $\operatorname{rad}(t)$ divides $\operatorname{ord}_q(\zeta_p^j b)$ and $\gcd\left(t, \frac{q-1}{\operatorname{ord}_q(\zeta_p^j b)}\right) = 1$. Hence every factor of the form $x^t - \zeta_p^j b$ is an irreducible factor of $x^n - 1$, which completes the first case.

Suppose now that $p = 2$, $q \equiv 3 \pmod 4$ and that for each odd prime factor p' of n we have that $\nu_{p'}(n) \leq \nu_{p'}(q-1)$. By the hypothesis, we know that

$$3 > \nu_2(n) > \nu_2(q-1) = 1,$$

so $n = 4m$ where m is an odd number such that $m|(q-1)$. In the case $m = 1$, the cyclotomic polynomial $\Phi_4(x) = x^2 + 1$ is irreducible, because -1 is not a square in $\mathbb{F}_q$. Thereby, we can suppose that $m \geq 3$ and by Proposition 2.4 we have

$$\Phi_n(x) = \Phi_{2m}(x^2) = \Phi_m(-x^2).$$

Now, since $\Omega(m) < \Omega(n)$, by induction hypothesis we know that every irreducible factor of $\Phi_m(x)$ is of the form $x^t - a$, so it satisfies the condition of Lemmas 2.1 and 2.3, and $x^{2t} - (-a)$ is a factor of $\Phi_n(x)$. We claim that it is an irreducible factor. In fact, since

$$\mathrm{ord}_q(-a) = \mathrm{lcm}(\mathrm{ord}_q(-1), \mathrm{ord}_q(a)),$$

then $\mathrm{ord}_q(-a)$ is an even number and then $\mathrm{rad}(2t)$ divides $\mathrm{ord}_q(-a)$. In addition, $\frac{q-1}{\mathrm{ord}_q(-a)}$ is an odd number, therefore $\gcd\left(2t, \frac{q-1}{\mathrm{ord}_q(-a)}\right) = 1$. Finally, since $t|m$ then $2t$ is not divisible by 4, thus by Lemma 2.1 it follows that $x^{2t} - (-a)$ is an irreducible factor of $x^n - 1$. □

Corollary 3.3. *Let n and q be as in Theorem 3.2 and set $m = \frac{n}{\gcd(n,q-1)}$ and $l = \frac{q-1}{\gcd(q-1,n)}$. Then*

(a) The factorization of $x^n - 1$ into irreducible factors of $\mathbb{F}_q[x]$ is

$$\prod_{t|m} \prod_{\substack{1 \leq u \leq \gcd(n,q-1) \\ \gcd(u,t)=1}} (x^t - \theta^{ul}).$$

(b) For each $t|m$, the number of irreducible factors of degree t is $\frac{\varphi(t)}{t} \cdot \gcd(n, q-1)$, where φ denotes the Euler Totient function and the total number of irreducible factors is

$$\gcd(n, q-1) \cdot \prod_{\substack{p|m \\ p \text{ prime}}} \left(1 + \nu_p(m)\frac{p-1}{p}\right).$$

Proof: (a) Let $x^t - a$ be an arbitrary irreducible factor of $x^n - 1$, and let p be a prime such that $p|t$. As a consequence of Lemma 2.1 we have that

$$\nu_p(\mathrm{ord}_q\, a) \geq 1 \text{ and } \nu_p(q-1) = \nu_p(\mathrm{ord}_q\, a).$$

In addition, as a consequence of Lemma 2.3 we have that

$$\nu_p(t) \leq \nu_p(n) \text{ and } \nu_p(\mathrm{ord}_q\, a) \leq \min\left\{\nu_p(q-1), \nu_p\left(\frac{n}{t}\right)\right\} \leq \nu_p(n) - \nu_p(t).$$

Therefore, for each prime p such that $\nu_p(t) \geq 1$, we have that $\nu_p(t) \leq \nu_p(n) - \nu_p(q-1)$, so we conclude that t divides $m = \frac{n}{\gcd(n,q-1)}$.

Now, note that $a = \theta^v$ for some $0 \leq v \leq q-1$, and since

$$\nu_p(q-1) = \nu_p(\mathrm{ord}_q\, a) = \nu_p(\mathrm{ord}_q\, \theta^v) = \nu_p\left(\frac{q-1}{\gcd(q-1, v)}\right) = \max\{0, \nu_p(q-1) - \nu_p(v)\},$$

we have that $\nu_p(v) = 0$, so it follows that $\gcd(t, v) = 1$.

Using that $\mathrm{ord}_q\, a | \frac{n}{t}$, we have that $a^{n/t} = \theta^{vn/t} = 1$, then vn/t is a multiple of the order of θ, i.e. $(q-1)|(vn/t)$, and therefore $\frac{q-1}{\gcd(q-1,n/t)} \Big| v$.

Since t is a divisor of $\frac{n}{\gcd(n,q-1)}$, we have that $\frac{q-1}{\gcd(q-1,n/t)}$ is a divisor of $l = \frac{q-1}{\gcd(q-1,n)}$. On the other hand, note that $\frac{q-1}{\gcd(q-1,n/t)}$ is a multiple of

$$\frac{q-1}{\gcd(q-1,n/m)} = \frac{q-1}{\gcd(q-1,\gcd(n,q-1))} = l.$$

therefore

$$\frac{q-1}{\gcd(q-1,n/t)} = \frac{q-1}{\gcd(q-1,n)}.$$

It follows that l divides v whatever is t, thus we can set $v = ul$, where $1 \leq u \leq \frac{q-1}{l} = \gcd(n, q-1)$.

Finally, since $\gcd(t,u)|\gcd(t,v) = 1$, we conclude that $\gcd(t,u) = 1$. This facts show that $x^n - 1$ divide $\prod_{t|m} \prod_{\substack{1\leq u\leq \gcd(n,q-1)\\ \gcd(u,t)=1}} (x^t - \theta^{ul})$.

Conversely, if t and u satisfy these conditions on the product, it can verified directly that $x^t - \theta^{ul}$ is an irreducible factor of $x^n - 1$, so *(a)* is proved.

(b) Let t be a divisor of $m = \frac{n}{\gcd(n,q-1)}$. Since

$$\operatorname{rad}(t)|\operatorname{rad}(n)|\gcd(q-1,n),$$

every prime that divides t also divides $\gcd(q-1,n)$. Let $p_1, p_2, \ldots, p_k$ be the list of primes that divide t. It follows that there exist $(1-\frac{1}{p_1})\cdot\gcd(q-1,n)$ numbers less or equal to $\gcd(q-1,n)$ that do not have any common factor with p_1, that there exist $(1-\frac{1}{p_1})(1-\frac{1}{p_2})\cdot\gcd(q-1,n)$ that do not have any common factor with p_1 and p_2, and inductively we conclude that there exist

$$\left(1-\frac{1}{p_1}\right)\left(1-\frac{1}{p_2}\right)\cdots\left(1-\frac{1}{p_k}\right)\cdot\gcd(q-1,n) = \frac{\varphi(t)}{t}\gcd(q-1,n)$$

numbers without any common factor with t. Finally the total number of irreducible factors of $x^n - 1$ is

$$\sum_{t|m}\frac{\varphi(t)}{t}\cdot\gcd(q-1,n).$$

Observe now that the function $\frac{\varphi(t)}{t}$ is a multiplicative function, therefore $\sum_{t|m}\frac{\varphi(t)}{t}$ is also multiplicative and thus it is enough to calculate this sum for powers of primes. In this case we have that

$$\sum_{d|p^k}\frac{\varphi(p^k)}{p^k} = 1 + k\left(1-\frac{1}{p}\right),$$

obtaining the part *(b)*. □

In the following theorem, we analyze the factorization of $x^n - 1$ in the case when $8|n$ and $q \equiv 3 \pmod 4$.

Theorem 3.4. *Let $\mathbb{F}_q$ be a finite field and n a positive integer such that*

(1) $q \equiv 3 \pmod 4$ and $8 \mid n$.
(2) $rad(n)|(q-1)$.

Then each irreducible factor of $x^n - 1$ in $\mathbb{F}_q[x]$ is of one of the following types:

(a) $x^t - a$ satisfying the hypothesis of Lemmas 2.1 and 2.3.
(b) $x^{2t} + (a+a^q)x^t + a^{q+1} \in \mathbb{F}_q[x]$, where $a \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$ and $x^t - a \in \mathbb{F}_{q^2}[x]$ satisfies the hypothesis of Lemmas 2.1 and 2.3 in $\mathbb{F}_{q^2}$.

Proof: Since $q^2 \equiv 1 \pmod 4$, it follows that the irreducible factors of $x^n - 1$ in $\mathbb{F}_{q^2}[x]$ are of the form $x^t - a$ and then they satisfy Lemmas 2.1 and 2.3 in $\mathbb{F}_{q^2}[x]$. Now, let $f(x) \in \mathbb{F}_q[x]$ be a monic irreducible factor of $x^n - 1$. From the fact that $f(x)$ is also a polynomial in $\mathbb{F}_{q^2}[x]$, we have two cases to consider:

i) If $f(x)$ is irreducible in $\mathbb{F}_{q^2}[x]$, then $f(x) = x^t - a$, where $a \in \mathbb{F}_q$, thus it satisfies the conditions of Lemmas 2.1 and 2.3 in $\mathbb{F}_q$.

ii) If $f(x)$ is not irreducible in $\mathbb{F}_{q^2}[x]$, then there exists $a \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q$ such that $(x^t - a) | f(x)$. Since $\sigma_q(f(x)) = f(x)$, where σ_q is the Frobenius automorphism

$$\begin{array}{rccc} \sigma_q : & \mathbb{F}_{q^2} & \to & \mathbb{F}_{q^2} \\ & b & \mapsto & b^q \end{array},$$

it follows that $\sigma_q(x^t - a) = x^t - a^q$ also divides $f(x)$. But $a \neq a^q$, therefore $(x^t - a)(x^t - a^q) | f(x)$. Finally, note that $(x^t - a)(x^t - a^q)$ is invariant by σ_q, hence $(x^t - a)(x^t - a^q) \in \mathbb{F}_q[x]$, and since $f(x)$ is irreducible, we conclude that $f(x) = x^{2t} - (a + a^q)x^t + a^{q+1}$. □

Remark 3.5. *With an equivalent proof of theorem above, but replacing the the hypotheses of the theorem by* $\mathrm{rad}(n) | (q^2 - 1)$*, or equivalently, every prime factor* p *of* n *satisfies that* $q \equiv \pm 1 \pmod p$*, it is possible to prove that every irreducible factor of* $x^n - 1$ *is a binomial or trinomial.*

Corollary 3.6. *Let* n *and* q *be as in Theorem 3.4. Let* α *be a generator of the cyclic group* $\mathbb{F}_{q^2}^*$ *satisfying* $\alpha^{q+1} = \theta$ *and put* $m = \frac{n}{\gcd(n, q^2-1)}$, $l_1 = \frac{q-1}{\gcd(q-1, n)}$, $l_2 = \frac{q^2-1}{\gcd(q^2-1, n)}$ *and* $r = \min\{\nu_2(n/2), \nu_2(q+1)\}$*. Then*

(a) The factorization of $x^n - 1$ *into irreducible factors of* $\mathbb{F}_q[x]$ *is*

$$\prod_{\substack{t | m \\ t \text{ odd}}} \prod_{\substack{1 \leq w \leq \gcd(n, q-1) \\ \gcd(w,t)=1}} (x^t - \theta^{w l_1}) \cdot \prod_{t | m} \prod_{u \in \mathcal{R}_t} (x^{2t} - (\alpha^{u l_2} + \alpha^{q u l_2}) x^t + \theta^{u l_2}),$$

where $\mathcal{R}_t$ *is the set*

$$\left\{ u \in \mathbb{N} \;\middle|\; \begin{array}{c} 1 \leq u \leq \gcd(n, q^2 - 1), \gcd(u, t) = 1 \\ 2^r \nmid u \text{ and } u < \{qu\}_{\gcd(n, q^2-1)} \end{array} \right\}$$

and $\{a\}_b$ *denotes the remainder of the division of* a *by* b*, i.e. it is the number* $0 \leq c < b$ *such that* $a \equiv c \pmod b$*.*

(b) For each t *odd with* $t | m$*, the number of irreducible binomials of degree* t *and* $2t$ *are* $\frac{\varphi(t)}{t} \cdot \gcd(n, q-1)$ *and* $\frac{\varphi(t)}{2t} \cdot \gcd(n, q-1)$ *respectively, and the number irreducible trinomials of degree* $2t$ *is*

$$\begin{cases} \frac{\varphi(t)}{t} \cdot 2^{r-1} \gcd(n, q-1) & \text{if } t \text{ is even} \\ \frac{\varphi(t)}{t} \cdot (2^{r-1} - 1) \gcd(n, q-1) & \text{if } t \text{ is odd.} \end{cases}$$

The total number of irreducible factors is

$$\gcd(n, q-1) \cdot \left(\frac{1}{2} + 2^{r-2}(2 + \nu_2(m)) \right) \cdot \prod_{\substack{p | m \\ p \text{ odd prime}}} \left(1 + \nu_p(m) \frac{p-1}{p} \right).$$

Proof: Since $\mathrm{rad}(n)|(q-1)$, we have that the only prime factor in common between n and $q+1$ is 2, so $\gcd(n/2, q+1) = 2^r$. In addition,

$$l_2 = \frac{q^2-1}{\gcd(q^2-1,n)} = \frac{q+1}{\gcd(q+1,n/2)}\frac{q-1}{\gcd(q-1,n)} = \frac{q+1}{2^r}l_1.$$

Now, we know by Corollary 3.3 that x^n-1 can be split in $\mathbb{F}_{q^2}[x]$ as

$$\prod_{t|m}\prod_{\substack{1\le u\le \gcd(n,q^2-1)\\ \gcd(u,t)=1}}(x^t-\alpha^{ul_2}).$$

Note that a factor $x^t-\alpha^{ul_2}$ is in $\mathbb{F}_q[x]$ if and only if α^{ul_2} is invarant by the Frobenius automorphism, i.e. $\alpha^{ul_2} = \alpha^{ul_2 q}$. This last equation is equivalent to saying that $(q^2-1)|(q-1)ul_2$, so $(q+1)|ul_2$. On the other hand

$$\begin{aligned}\gcd(q+1,l_2) &= \gcd\left(q+1, \frac{q+1}{\gcd(n/2,q+1)}\cdot\frac{q-1}{\gcd(n,q-1)}\right)\\ &= \gcd\left(q+1, \frac{q+1}{\gcd(n/2,q+1)}\right)\\ &= \frac{q+1}{\gcd(n/2,q+1)} = \frac{q+1}{2^r},\end{aligned}$$

so $x^t-\alpha^{ul_2}$ is in $\mathbb{F}_q[x]$ if and only if $2^r|u$. Thus, we can write $u = 2^r w$, and since $\gcd(u,t)=1$, it follows that t is odd and $\gcd(w,t)=1$. Moreover, from the fact that

$$ul_2 = 2^r w\frac{q+1}{2^r}l_1 = (q+1)wl_1,$$

we have that $x^t-\alpha^{ul_2} = x^t-\alpha^{(q+1)wl_1} = x^t-\theta^{wl_1}$, and these conditions determine every irreducible factor of x^n-1 that is binomial in $\mathbb{F}_q[x]$.

For the values of u such that $2^r \nmid u$, we have that $x^t-\alpha^{ul_2}\notin\mathbb{F}_q[x]$ and then

$$(x^t-\alpha^{ul_2})(x^t-\alpha^{qul_2}) = x^{2t}-(\alpha^{ul_2}+\alpha^{qul_2})x^t+\theta^{ul_2}$$

is an irreducible trinomial of $\mathbb{F}_q[x]$. Thus, each pair $(ul_2, \{uql_2\}_{q^2-1})$, or equivalently each pair $(u, \{qu\}_{\gcd(n,q^2-1)})$, generates an irreducible trinomial. It follows that each irreducible trinomial is generated by a unique pair in the set

$$\left\{(u,v)\;\middle|\;\begin{array}{l}1\le u\le \gcd(n,q^2-1), \gcd(u,t)=1\\ v=\{qu\}_{\gcd(n,q^2-1)}, 2^r\nmid u \text{ and } \ u<v\end{array}\right\}$$

and this concludes the part *(a)*.

An important fact to emphasize is that, if $\alpha^{ul_2}+\alpha^{qul_2}=0$, then the "trinomial" $x^{2t}-(\alpha^{ul_2}+\alpha^{qul_2})x^t+\theta^{ul_2}$ is really a binomial of degree $2t$ with t odd.

Now, by Corollary 3.3, we know that for every t divisor of m, the number of irreducible binomials in $\mathbb{F}_{q^2}[x]$ is

$$\frac{\varphi(t)}{t}\gcd(n,q^2-1) = \frac{\varphi(t)}{t}2^r\gcd(n,q-1).$$

If t is even, then u is odd and there are not binomials of degree t in $\mathbb{F}_{q^2}[x]$ that are also in $\mathbb{F}_q[x]$. Thereby, for every binomial in $\mathbb{F}_{q^2}[x]$, there exists a unique binomials in $\mathbb{F}_{q^2}[x]$, such that its product generates an irreducible trinomial in $\mathbb{F}_q[x]$. So the number of irreducible trinomials of degree $2t$ is $\frac{\varphi(t)}{t}2^{r-1}\gcd(n,q-1)$.

Now, if t is odd there exist $\frac{\varphi(t)}{t}\gcd(n, q-1)$ irreducible binomials in $\mathbb{F}_{q^2}[x]$ than are also in $\mathbb{F}_q[x]$, and $\frac{\varphi(2t)}{2t}\gcd(n, q-1)$ reducible binomials in $\mathbb{F}_{q^2}[x]$ that are irreducible binomial in $\mathbb{F}_q[x]$. Therefore, there are

$$\frac{\varphi(t)}{t}\gcd(n, q^2-1) - 2\frac{\varphi(t)}{t}\gcd(n, q-1) = \frac{\varphi(t)}{t}(2^r - 2)\gcd(n, q-1)$$

binomials in $\mathbb{F}_{q^2}[x]$ than are not in $\mathbb{F}_q[x]$, so the number of irreducible trinomials of degree $2t$ is $\frac{\varphi(t)}{2t}(2^r-2)\gcd(n, q-1)$.

Finally, the total number of irreducible factors is

$$\begin{aligned}
&= \sum_{\substack{t|m \\ t \text{ odd}}} \frac{\varphi(t)}{2t}(2^r+1)\cdot\gcd(q-1,n) + \sum_{\substack{t|m \\ t \text{ even}}} \frac{\varphi(t)}{t}2^{r-1}\cdot\gcd(q-1,n) \\
&= \sum_{\substack{t|m \\ t \text{ odd}}} \frac{\varphi(t)}{2t}\cdot\gcd(q-1,n) + \sum_{\substack{t|m \\ t}} \frac{\varphi(t)}{t}2^{r-1}\cdot\gcd(q-1,n) \\
&= \gcd(n,q-1)\left(\frac{1}{2}\cdot\prod_{\substack{p|m \\ p \text{ odd prime}}}\left(1+\nu_p(m)\frac{p-1}{p}\right) + 2^{r-1}\cdot\prod_{\substack{p|m \\ p \text{ prime}}}\left(1+\nu_p(m)\frac{p-1}{p}\right)\right) \\
&= \gcd(n,q-1)\cdot\left(\frac{1}{2}+2^{r-2}(2+\nu_2(m))\right)\cdot\prod_{\substack{p|m \\ p \text{ odd prime}}}\left(1+\nu_p(m)\frac{p-1}{p}\right),
\end{aligned}$$

as we wanted to prove. □

Acknowledgements. We would like to thank anonymous referees for the comments about the first version of this article, in particular, by propose the question about the converse of Theorem 3.2.

References

[1] Berlekamp, E.R., *Bit-serial Reed-Solomon encoders*. IEEE Trans. Info. Theory **28** (1982) 869-874.

[2] Blake, I. F., Gao, S., Mullin, R. C., *Explicit factorization of $x^{2^k}+1$ over $\mathbb{F}_p$ with $p \equiv 3 \pmod 4$*, Appl. Algebra Engrg. Comm. Comput. **4** (1993) 89-94.

[3] Chen, B., Li, L., Tuerhong, R., *Explicit factorization of $x^{2^m p^n} - 1$ over a finite field*. Finite fields and Their Applications **24** (2013) 95-104.

[4] Golomb, S. W., *Shift Register Sequences* Holden-Day, Inc. 1967.

[5] Lidl, R., Niederreiter, H., *Finite Fields*. Encyclopedia of Mathematics and Its Applications, Vol 20, Addison-Wesley 1983.

[6] Van Lint J.H., *Introduction to Coding Theory (3rd ed)*, Graduate Texts in Mathematics 86, Springer Verlag, 1998.

[7] Fitzgerald R.W., Yucas J.L., *Explicit factorization of cyclotomic and Dickson polynomials over finite fields*. Arithmetic of Finite Fields. Lecture Notes in Computer Science, vol. 4547, pp. 1-10. Springer, Berlin (2007).

[8] Meyn H., *Factorization of the cyclotomic polynomials $x^{2^n}+1$ over finite fields*. Finite Fields Appl. 2, (1996) 439-442.

[9] Wang, L.; Wang, Q., *On explicit factors of cyclotomic polynomials over finite fields*. Des. Codes Cryptogr. 63 (2012), no. 1, 87-104.

Departamento de Matemática, Universidade Federal de Minas Gerais, UFMG, Belo Horizonte, MG, 30123-970, Brazil,
E-mail address: fbrocher@mat.ufmg.br
E-mail address: carmita@mat.ufmg.br
E-mail address: liubatista20@gmail.com